VERTICAL EXAGGERATION X 2

GEOLOGIC MAP OF THE ST. JOE QUADRANGLE, SEARCY AND MARION COUNTIES, ARKANSAS

Mark R. Hudson and Kenzie J. Turner

St. Joe quadrangle, Searcy and Marion Counties, Arkansas: U.S. Geological Survey Scientific Investigations Map 3074,

STRUCTURAL GEOLOGY

may contain medium- to light-gray, fetid septarian concretions as large as 2 ft in diameter. The Fayetteville Shale is susceptible to landslides.

Batesville Sandstone (Upper Mississippian, Chesterian)—Fine-grained to

very fine grained, light- to medium-brown, calcite-cemented sandstone with

sparse interbedded limestone. Thin to medium beds are typically parallel

laminated; low-angle crossbeds common in upper part of unit. Sandstone

commonly contains burrows on bedding plane surfaces. Sandstone breaks

into thin flat blocks. One or more discontinuous, 1- to 3-ft-thick, medium-

to dark-gray, fetid, fossiliferous limestone beds are locally interbedded with

sandstone and limestone beds may contain 2- to 10-mm-diameter oxidized

pyrite framboids that weather to reddish-brown spheres. Where stripped of

overlying Fayetteville Shale, top of Batesville forms topographic flats. Unit

Member. The Boone Formation is a common host of caves and sinkholes.

hosts sinkholes formed by collapse into dissolution cavities in underlying

Boone Formation (Upper to Lower Mississippian)—Limestone and cherty limestone of main body that grade into the basal St. Joe Limestone

Main body (Upper to Lower Mississippian, Meramecian to Osagean)-Medium- to thick-bedded, chert-bearing bioclastic limestone. Limestone is

light to medium gray on fresh surfaces and generally coarsely crystalline

present in upper one-third of unit. Beds are typically parallel planar to wavy, but channel fills are locally present in lower part of unit. Chert

content varies vertically and laterally within Boone and forms lenticular to

produce abundant float of white-weathered chert on hillslopes. Chert in

uppermost part of unit contains prominent brachiopod molds. Thickness

Kinderhookian)—Thin and wavy bedded bioclastic limestone containing ubiquitous 3- to 6-mm-wide crinoid fragments in a finely crystalline matrix.

Limestone is commonly pink to red on fresh surfaces due to hematite in

matrix, but color varies depending on hematite concentration. Lenticular

red to pink chert nodules are locally present in upper part. Contact with

Limestone Member contains greenish-gray shale interbeds. Base of St. Joe

unconformity that variably truncates Fernvale Limestone, Plattin Limestone, St. Peter Sandstone, or upper part of the Everton Formation in various

overlying main body of Boone Formation is gradational and marked by increase in chert and thicker beds upward. Middle to lower part of St. Joe

is a 0.3- to 8-ft-thick, very fine to fine-grained, moderately sorted, tan

sandstone containing phosphate pebbles. Base of unit is an important

Fernvale Limestone (Upper Ordovician)—Thick- to massive-bedded, coarsely

crystalline bioclastic limestone. Limestone is light gray to pink on fresh

shaped crinoid ossicles in a coarsely crystalline matrix. Mapped undivided

with Fernvale Limestone is a 5-ft-thick fissile to massive black to blue-green

shale with 1 to 2 in. black phosphate pebbles, which is assigned to Cason

Shale (Upper Ordovician) where it was observed beneath a cliff of St. Joe

Limestone Member adjacent to Mill Creek (NE1/4, NE1/4, Sec. 18, T. 16

fine-grained, dense limestone locally interbedded with calcarenite. Charac-

light-gray, fine-grained to very fine grained, limey to dolomitic calcarenite as

thick as 10 ft; calcarenite has fetid odor when freshly broken. Thickness

ranges from 40 ft in southwestern part of area to nothing in northwestern

parts due to pre-Mississippian erosion beneath unconformity at base of St.

sandstone interbedded with blue-green siltstone, shale, and sparse dolos-

tone. Sandstone is calcite-cemented arenite containing well-rounded quartz

grains. Upper part of unit, as thick as 25 ft, is sandstone interbedded with

blue-green fissile shale, thin-bedded siltstone, and sparse 0.5- to 3-ft-thick

dolostone interval poorly exposed, but is locally marked by zone of slumped

sandstone blocks on hill slopes. Main part of unit contains a sequence of 1to 2-ft-thick sandstone beds overlying a massive basal sandstone ledge as

beds of brown-gray, fine crystalline dolostone; shale, siltstone, and

thick as 20 ft. Bioturbation is common in most sandstone beds and

includes cylindrical burrows (Skolithos) perpendicular to bedding that

contains inclined fractures, and discordantly overlies laminated sandy

Everton Formation (Middle Ordovician)—Interbedded dolostone, limestone,

and sandstone sequence, divided into upper and lower parts. Unit is

Upper part—Interbedded dolostone, sandstone, and limestone. Limestone is

finely crystalline, light gray, thick bedded, and locally fossiliferous, and is

present in intervals as thick as 20 ft. Dolostone is light to dark gray, finely

to medium crystalline, laminated, and medium to thick bedded commonly

well-sorted, well-rounded, fine to medium quartz grains. Sandstone is light

limestone. Limestone interbedded with limy sandstone and lesser dolostone

tan to white in planar, medium to thick beds and is variably cemented by

containing sandstone stringers. Sandstone is arenite composed of

dolomite or calcite; sandstone locally grades into sandy dolostone or

at top of unit comprises Jasper Member of Glick and Frezon (1953). Jasper Member is up to 40 ft thick but has been removed in northwestern

part of quadrangle due to pre-Mississippian erosion. Dolostone and

part of Everton. Base of upper part of Everton is marked by base of

dolomitic sandstone underlying Jasper Member make up most of upper

quartz arenite composed of well-sorted, well-rounded, fine to medium

Everton Formation is 180–200 ft thick

Newton Sandstone Member (McKnight, 1935), a planar, medium-bedded

quartz grains. Newton Sandstone Member is 5–10 ft thick. Upper part of

Lower part—Interbedded dolostone, sandstone, and limestone. Limestone is

hemispheres as large as 1 ft in diameter. Dolostone, present throughout

unit, is light gray to grayish brown, finely to medium crystalline, thin to

tolites. Sandstone is light-tan to white quartz arenite, composed of

well-sorted, well-rounded, fine to medium grains, variably cemented by

dolomite or calcite. Sandstone intervals, composed of planar, thin to

thick bedded, and commonly laminated, and it commonly contains stroma-

medium beds as thick as 6 ft, are present throughout unit. Sandstone also

forms lenses and stringers within limestone and dolostone intervals. Light-

to dark-brownish-gray dolostone and sandy dolostone interspersed with

sandstone are predominant in lower part of unit. Collapse breccia with

clasts of planar laminated dolomite and dolomitic sandstone is present.

Unit is 250 ft thick where fully exposed near Mt. Hersey to the west-

only upper 60 ft is exposed within St. Joe quadrangle

section only

present in its isolated western outcrops.

Lower Ordovician rocks, undivided—Sedimentary rocks shown on cross

STRATIGRAPHY

on what is now the southern margin of the North American continent. Provincial series

The Middle Ordovician Everton Formation (Oe) is a heterogeneous sandstone and

tidal flat depositional environments. The Everton Formation is discontinuously overlain by

eastern part of the Mill Creek graben, by the Upper Ordovician Fernvale Limestone (Of) and

Cason Shale (McKnight, 1935). Due to its limited thickness and distribution, Cason Shale

was mapped undivided with Fernvale Limestone where it was observed west of the town of

Batesville, Arkansas, Craig (1975) demonstrated that only its basal Upper Ordovician part is

The base of the Mississippian section is a major unconformity that progressively cuts

St. Joe. Whereas Cason Shale includes Silurian facies in its type area to the east near

deeper into the underlying Ordovician section towards the northwestern part of the

quadrangle. The Mississippian Boone Formation (Mb) is the most widespread unit at the

St. Joe Limestone Member (Mbs) is persistent throughout much of northern Arkansas

(McKnight, 1935). This sandstone was probably deposited as a transgressive lag during

sea-level rise in Late Devonian time (Horner and Craig, 1984). The contact of the main

body of the Boone Formation with the St. Joe Limestone Member is gradational and is

based on the change to thin bedding and the generally chert-free lithology of the St. Joe Limestone Member. Near the top of the formation an oolite interval as thick as 3 ft is

common. Braden and Ausbrooks (2003a) also recognized this facies in areas to the south

and correlated this interval with the Short Creek Oolite Member of the Boone Formation,

which is present in Missouri and Kansas (McKnight and Fischer, 1970). Black shales of the

Fayetteville Shale (Mf) represent deposition within a southward-deepening muddy shelf that

the Mississippian (Sutherland, 1988). Typical thin-bedded, rippled, fine- to very fine grained

sandstones and siltstones of Cane Hill Member (Phc) of the Hale Formation were probably

deposited in tidal flat environments (Manger and Sutherland, 1992). The quadrangle lies at

the northeastern margin of the distribution of thick Prairie Grove Member (Phg) of the Hale

Formation (Sutherland, 1988). The Prairie Grove Member in the study area is represented

by indistinct planar- or cross-bedded, calcite-cemented sandstone deposited in a shoreface

Grove Member and represents deposition in an open-marine platform setting as sea level

continued to rise. Pennsylvanian sandstone and shale of the upper part of the Bloyd

Formation (Pbu) were originally called Winslow Formation by Purdue and Miser (1916),

with the basal Greenland Sandstone Member of the Atoka Formation (Henbest, 1953)

concluded that the cliff-forming sandstone was a time-equivalent unit with the Woolsey

Member of the Bloyd Formation farther west and designated it with the informal term

of the Bloyd Formation represents a period of erosion prior to deposition of the middle

Fault—Dashed where approximately located

shown in cross sections only

Strike and dip of bedding

contour interval 50 ft

top of Boone Formation

Normal fault—Dashed where approximately located,

Strike-slip fault, right-lateral offset—Dashed where

Bloyd sandstone by terrestrial braided streams.

----- Contact

— Anticline Syncline

CONVERSION FACTORS

inches (in.)

feet (ft)

Monocline

★ Cason Shale locality

centimeters (cm

meters (m)

representing the prominent cliff-forming crossbedded sandstone. Zachry (1977), however,

"middle Bloyd sandstone." The unconformable contact between the upper and lower parts

MAP EXPLANATION

dotted where concealed, bar and ball on downthrown block

approximately located, arrows show relative motion

⊗ ⊙ **Fault movement**—Away from viewer (left), toward viewer (right),

—900— Line of equal elevation showing base of Boone Formation,

Control point showing elevation in feet on base or

environment. The lower part of the Bloyd Formation (PbI) conformably overlies the Prairie

Initial Morrowan deposition reflects sea-level rise following a sea-level drop at the end of

shoaled upward during deposition of the Pitkin Limestone (Mp) (Handford, 1985).

surface within the quadrangle. The phosphate-nodule-bearing sandstone at the base of the

carbonate unit that Suhm (1974) interpreted to have been deposited in barrier island and

the Middle Ordovician St. Peter Sandstone (Osp) and Plattin Limestone (Opl) and, in the

terms for Pennsylvanian and Mississippian units are from McFarland (1988).

The quadrangle preserves a 1,300-ft-thick record of early and late Paleozoic deposition

southwest on the Western Grove quadrangle (Glick and Frezon, 1953) but

restricted to upper part of unit and is interbedded with dolostone and

sandstone. Limestone is light gray, finely crystalline, thin to medium

bedded, commonly laminated, and contains sparse stromatolite

weather to distinctive straw-like forms. Basal sandstone is crossbedded,

dolomite in upper part of Everton Formation at local recesses. Unit thins

to nothing to northwest due to pre-Mississippian erosion beneath unconfor-

mity at base of St. Joe Limestone Member of Boone Formation. Thickness

teristically, dense, very fine grained, medium-gray limestone that breaks

with conchoidal fracture and weathers to very light gray, thin, tabular

blocks. Locally in upper parts there are interbeds of laminated, tan to

N., R. 17 W.). Unit is exposed in eastern Mill Creek graben where it is as

Plattin Limestone (Middle Ordovician)—Thin- to medium-planar-bedded,

Joe Limestone Member of Boone Formation

Osp St. Peter Sandstone (Middle Ordovician)—Very fine grained, grayish-yellow

surfaces and contains abundant 3- to 10-mm-wide cylindrical to barrel-

parts of quadrangle. Total thickness of member is 20–50 ft

St. Joe Limestone Member (Lower Mississippian, Osagean to

anastomosing lenses. Chert-rich horizons are generally poorly exposed but

with interspersed crinoid ossicles. A 1- to 3-ft-thick bed of oolitic limestone

is common in upper part of unit. Beds of dense, fine-grained limestone are

sandstone; limestone beds contain crinoids and brachiopods. Both

Thickness 130–160 ft

Boone Formation. Thickness 40–60 ft

Total thickness is 380–405 ft

thick as 20 ft

ranges from 0 to 70 ft

300–400 ft thick

Rocks within the map area were deformed by a system of faults and folds. Structure contours on the base of the Boone Formation illustrate the location of structures and their vertical offset. The structure contours conform to elevations at 328 control points located at both lower and upper contacts of the Boone Formation, as well as other information limiting maximum or minimum elevations of these contacts. A 390-ft thickness for the Boone Formation (Mb) (including the St. Joe Limestone Member (Mbs)) was used to project the elevation of the basal contact from points on the upper contact. This average thickness was based on thicknesses from five stratigraphic sections farther west near the Buffalo River (Hudson, 1998).

Several regionally extensive fault systems intersect within the St. Joe quadrangle. We

map faults that correspond to those previously named by McKnight (1935) as the St. Joe, Mill Creek, Pilot Mountain, Tomahawk, and South Tomahawk faults, although our fault traces differ in detail, and we also map several additional faults. The Mill Creek graben in the southwestern part of the quadrangle is part of a graben system that extends about 15 miles farther to the west (Purdue and Miser, 1916; Hudson and Murray, 2003; Hudson and others, 2006). Bounding the south side of the graben, the Mill Creek fault dies out eastward at its intersection with the northeast-striking Confederate fault, which we project into the quadrangle from areas farther southwest (Braden and Ausbrooks, 2003a, 2003b). Structure contours highlight a gentle southeast dip of strata in a broad flank northwest of the Confederate fault. This dipping block was part of the St. Joe monocline as named by McKnight (1935), but we abandon this as a separate structural feature and instead link it to growth of the Confederate fault. Cross section A-A' demonstrates that about 200 feet down-to-thesoutheast throw on the Confederate fault is required in addition to the southeast dip of strata to provide sufficient structural relief for the typical thickness of Boone Formation (Mb) between outcrops of its base and top. Bounding the north side of the Mill Creek graben, the St. Joe fault continues northeastward from its intersection with the Confederate fault until it overlaps the Davy Crockett fault at the southeastern border of Sec. 8, T. 16 N., R. 17 W. The Davy Crockett fault includes the northeast-striking segment of the Tomahawk fault of McKnight (1935), but we rename this fault segment because our mapping demonstrates that it continues to the northeast past McKnight's west-northwest striking segment of the Tomahawk fault, which we retain as a name. Similar to the Tomahawk fault, the west-

northwest-striking South Tomahawk and Pilot Mountain faults extend from beyond the east

edge of the quadrangle to intersect the Davy Crockett fault at their west ends. The West

Mountain fault aligns with, and may be a westward continuation of, the Pilot Mountain fault across the Davy Crockett fault. Whereas the vertical offset of structures can be estimated from the elevation difference of formation contacts across the structures, their lateral offset is difficult to measure due to the lack of appropriate markers. Fault data collected from mostly small-scale faults at numerous sites within the quadrangle (fig. 3) indicate the typical orientations and slip senses for faults (fig. 4A) and give insight into the causative stresses responsible for deformation. Most of these faults are small displacement cataclastic faults, deformation bands (Aydin and Johnson, 1978), preferentially developed in sandstones of the St. Peter Sandstone (Osp) and Everton Formation (Oe). Those faults whose slip senses could be determined are predominantly normal or strike slip (fig. 4B). Normal faults strike from east-northeast to east-southeast and dip moderately to the north or south. Among strike-slip faults, those with right-lateral sense mostly strike northeast and those with left-lateral sense have strikes ranging from east-southeast to south-southeast. Paleostress inversions for these normal and strike-slip faults, using the method of Angelier (1990), demonstrate that they formed in a stress regime with a least principal stress axis oriented south to south-southeast and near horizontal and with intermediate and maximum principal stress axes flipping between near vertical and east-northeast and near horizontal (fig. 4B). Similar coordinated movements among normal and strike-slip faults have been documented elsewhere in the southern Ozark dome, and their north-south extension accommodated late Paleozoic flexure of the southern continental margin as it was loading by the adjacent Ouachita orogenic belt (Hudson, 2001). From these data we interpret that (1) northeast-striking Confederate and Davy Crockett faults have components of right-lateral slip as well as down-to-the-southeast throw, and (2)

Crockett (cross section B-B') and Pilot Mountain faults. Two east-west-trending open anticlines and a syncline lie directly north of the St. Joe fault. A broad, east-southwesttrending dome is present in the northeastern part of the quadrangle. Away from folds, dips of bedding are low and variable in direction. These dispersed attitudes are attributed in part to local subsidence caused by karst dissolution within the abundant limestone and dolostone Joints measured within the map area (695 total) are near vertical and distributed in several sets (fig. 4). The dominant set strikes north. Less prominent joint sets strike

east- to southeast-striking faults like those bounding the Mill Creek graben have normal slip.

monocline on the eastern edge of the quadrangle has about 150 ft of relief and is expressed

underlies St. Joe and Pilot Mountains with a northern limb upturned adjacent to the Davy

by consistent $5-10^{\circ}$ north to north-northeast dips. Just to the north, a synclinal axis

The structure contours illustrate several folds and broad domes. The Baker Spring

northeast and west-northwest. Joint planes within limestone and dolostone formations, such as the Boone (Mb) and Everton Formations (Oe), are commonly enlarged due to dissolution. **ECONOMIC GEOLOGY**

McKnight (1935) described lead and zinc ores from several mines in the St. Joe and Tomahawk Creek mining districts located on the northern, upthrown side of the St. Joe and Davy Crockett faults within the St. Joe quadrangle (fig. 3). The dominant ore mined from local prospects consists of sphalerite (ZnS) and zinc carbonate and minor copper in the form of malachite. Dolomitic and limestone layers of the Everton Formation (Oe) host most prospects. Silicification was commonly associated with Pb-Zn mineralization (McKnight, 1935) and was noted at several sites during this field investigation (fig. 3) defining an elongate area mostly north of the St. Joe and Davy Crockett faults. Within the Boone Formation (Mb), silica was precipitated as fracture coatings of drusy quartz having euhedral crystals tips as wide as $\frac{1}{2}$ in. (fig. 5A). Additionally, within the Boone Formation and its basal St. Joe Limestone Member, several elongate bodies of brecciated chert cemented by microcrystalline quartz (fig. 5B) were noted, standing 3 to 20 ft high as erosion-resistant ridges. These masses probably formed by dissolution of limestone and collapse of chert within the Boone Formation followed by silicification during flow of mineralizing fluids. Enhanced silicic cementation of sandstone beds was also noted within the Ordovician St. Peter Sandstone (Osp) and Everton Formation (Oe) and at one location within the Pennsylvanian Cane Hill Member (Phc) of the Hale Formation.

ACKNOWLEDGMENTS

This work was conducted in a cooperative project between the U.S. Geological Survey and the U.S. National Park Service. Special thanks go to the many private owners who allowed access onto their land during this study.

REFERENCES CITED

Angelier, Jacque, 1990, Inversion of field data in fault tectonics to obtain the regional stress,

III, A new rapid direct inversion method by analytical means: Geophysical Journal International, v. 103, p. 363–376. Aydin, A., and Johnson, A.M., 1978, Development of faults as zone of deformation bands and as slip surfaces in sandstone: Pure and Applied Geophysics, v. 116, p. 931–942. Braden, A.K., and Ausbrooks, S.M., 2003a, Geologic map of the Eula quadrangle, Newton and Searcy Counties, Arkansas: Arkansas Geological Commission Digital Geologic Quadrangle Map DGM-AR-00269, scale 1:24,000. Braden, A.K., and Ausbrooks, S.M., 2003b, Geologic map of the Snowball quadrangle,

Searcy County, Arkansas: Arkansas Geological Commission Digital Geologic Quadrangle Map DGM-AR-00800, scale 1:24,000. Craig, W.W., 1975, Stratigraphy and conodont faunas of the Cason Shale and the Kimmswick and Fernvale Limestones of northern Arkansas, in Headrick, K.N., and Wise, O.A., eds., Contributions of the geology of the Arkansas Ozarks, Arkansas Geological

Commission, p. 61–95. Glick, E.E., and Frezon, S.E., 1953, Lithologic character of the St. Peter Sandstone and Everton Formation in the Buffalo River Valley, Newton County, Arkansas: U.S. Geological Survey Circular 249, 39 p.

Handford, C.R., 1985, Facies and bedding sequences in shelf-storm-deposited carbonates— Fayetteville Shale and Pitkin Limestone (Mississippian), Arkansas: Journal of Sedimentary Petrology, v. 56, p. 123-137. Henbest, L.G., 1953, Morrow Group and lower Atoka Formation of Arkansas: American Association of Petroleum Geologists Bulletin, v. 37, p. 1935–1953. Horner, G.J., and Craig, W.W., 1984, The Sylamore Sandstone of north-central Arkansas, with emphasis on the origin of its phosphate, in McFarland, J.D., III, ed., Contributions to the geology of Arkansas, v. 2: Arkansas Geological Commission, p. 51–85.

Hudson, M.R., 1998, Geologic map of parts of the Jasper, Hasty, Ponca, Gaither, and Harrison quadrangles in and adjacent to Buffalo National River, northwestern Arkansas: U.S. Geological Survey Open-File Report 98–116, scale 1:24,000. Hudson, M.R., 2001, Coordinated strike-slip and normal faulting in the southern Ozark dome of northern Arkansas: Deformation in a late Paleozoic foreland: Geology, v. 28, p. 511–514.

Hudson, M.R., Murray, K.E., and Pezzutti, D., 2001, Geologic map of the Jasper quadrangle, Newton and Boone Counties, Arkansas: U.S. Geological Survey Miscellaneous Field Studies Map MF-2356, scale 1:24,000. Hudson, M.R., and Murray, K.E., 2003, Geologic map of the Ponca quadrangle, Newton, Boone, and Carroll Counties, Arkansas: U.S. Geological Survey Miscellaneous Field Studies Map MF-2412, scale 1:24,000.

Hudson, M.R., and Murray, K.E., 2004, Geologic map of the Hasty quadrangle, Newton and Boone Counties, Arkansas: U.S. Geological Survey Scientific Investigations Map 2847, scale 1:24,000, http://pubs.usgs.gov/sim/2004/2847/. Hudson, M.R., Turner, K.J., and Repetski, J.E., 2006, Geologic map of the Western Grove Quadrangle, Northwestern Arkansas: U.S. Geological Survey Scientific Investigations Map 2921, 1:24,000 scale, http://pubs.usgs.gov/sim/2006/2921/. Hudson, M.R., and Turner, K.J., 2007, Geologic map of the Boxley Quadrangle, Newton and Madison Counties, Arkansas: U.S. Geological Survey Scientific Investigations Map

Manger, W.L., and Sutherland, P.K., 1992, Analysis of sections presumed to be complete across the Mississippian-Pennsylvanian boundary, southern mid-continent, in Sutherland, P.K., and Manger, W.L., eds., Recent advances in middle Carboniferous biostratigraphy—A symposium: Oklahoma Geological Survey Circular 94, p. 69–79. McFarland, J.D., III, 1988, The Paleozoic rocks of the Ponca region, Buffalo National River, Arkansas, in Hayward, O.T., ed., Centennial Field Trip Guide, v. 4: Geological Society of America, p. 207–210.


2991, 1:24,000 scale, http://pubs.usgs.gov/sim/2991/.

100, p. 1787–1802.

McKnight, E.T., 1935, Zinc and lead deposits of northern Arkansas: U.S. Geological Survey Bulletin 853, 311 p. McKnight, E.T., and Fischer, R.P., 1970, Geology and ore deposits of the Picher field, Oklahoma and Kansas: U.S. Geological Survey Professional Paper 588, 165 p. Purdue, A.H., and Miser, H.D., 1916, Descriptions of the Eureka Springs and Harrison quadrangles: U.S. Geological Survey Atlas, Folio 202, scale 1:125,000. Suhm, R.A., 1974, Stratigraphy of Everton Formation (early Medial Ordovician), northern Arkansas: American Association of Petroleum Geologists Bulletin, v. 58, p. 685–707. Sutherland, P.K., 1988, Late Mississippian and Pennsylvanian depositional history in the


Zachry, D.L., 1977, Stratigraphy of middle and upper Bloyd strata (Pennsylvanian, Morrowan), northwestern Arkansas, in Sutherland, P.K., and Manger, W.L., eds., Upper Chesterian-Morrowan stratigraphy and the Mississippian-Pennsylvanian boundary in northeastern Oklahoma and northwestern Arkansas: Oklahoma Geological Survey Guidebook 18, p. 61–66.

Arkoma basin area, Oklahoma and Arkansas: Geological Society of America Bulletin, v.




SCIENTIFIC INVESTIGATIONS MAP 3074

Figure 1. Location of study area within northern Arkansas, adjacent to the western part of Buffalo National River. Other U.S. Geological Survey geologic maps in area (Hudson, 1998; Hudson and others, 2001, 2006; Hudson and Murray, 2003, 2004; Hudson and Turner, 2007) are listed by publication number. Lower regional map illustrates geological and selected physiographic provinces of Arkansas and adjacent areas.



EXPLANATION Silicification site Fault site Mine or prospect site **Fault**—Dashed where approximately located, dotted where concealed

Figure 3. Shaded-relief digital elevation map of quadrangle with locations of Pb-Zn mines and prospects, structural features, and sites where faults (squares) and silicification (triangles) were observed. Where determined, slip senses for faults at sites are indicated as normal, N (rake 90° to 60°); oblique, O (rake 59° to 30°); and strike-slip, S (rake 29° to 0°). Sites with silicification define an elongate area (pink) centered mostly north of the St. Joe and Davy Crockett faults. Silicification was classified as Q, coarse and microcrystalline quartz crystals; B, silica-cemented chert breccia; or C, enhanced silica cementation in sandstone. Mine and prospect locations are estimated from descriptions in McKnight (1935).





Boone Formation (pencil is about 5 in. long), and (B) chert breccia mass (hammer handle is about 12 in. long). Photographs by M.R. Hudson and K.J. Turner, 2008.

Any use of trade names is for descriptive purposes only and does not imply endorsement by the U.S. Government This map was produced on request, directly from digital files, on an electronic plotter For sale by U.S. Geological Survey Information Services Box 25286, Federal Center, Denver, CO 80225 1-888-ASK-USGS ArcInfo coverages and a PDF for this map are available at http://pubs.usgs.gov/sim/3074/ Hudson, M.R., and Turner, K.J., 2009, Geologic map of the